

Name _____

Honor Chemistry

__/__/__

Energy

Energy is the ability to do work or produce heat. Energy can be classified as either **kinetic energy** (energy of motion) or **potential energy** (energy due to position or composition).

One step in understanding energy is realizing the difference between **heat** and **temperature**. Temperature is a measure of the average kinetic energy of the particles in a substance. Heat is the flow of energy due to a temperature difference. Heat travels from a higher temperature object to a lower temperature object until both objects reach the same temperature. The **law of conservation of energy** states that energy cannot be created or destroyed. So in all heat transfers one object is **endothermic** (absorb heat) and the other is **exothermic** (release heat).

In chemistry we measure heat using two different units: calories and joules. A **calorie** is defined as the amount of energy needed to raise the temperature of one gram of water one degree Celsius. The “calorie” you are most familiar with is used to measure the energy content of food. This “calorie” is actually a kilocalorie (1000 calories) and is written with a capital C to distinguish it from the calorie used in chemistry.

The Calories listed for this candy bar are actually kilocalories.

Nutrition Facts		Amount/Serving	%DV*	Amount/Serving	%DV*
Serving Size 1 bar		Total Fat 11 g	17%	Total Carb. 28 g	9%
Calories 230		Sat. Fat 8 g	40%	Dietary Fiber <1 g	3%
Fat Cal. 100		Cholest. <5 mg	2%	Sugars 22 g	
*Percent Daily Values (DV) are based on a 2,000 calorie diet.		Sodium 135 mg	6%	Protein 4 g	
		Vitamin A 0% • Vitamin C 0% • Calcium 4% • Iron 2%			

The other unit used to measure heat is the **joule**. The joule is the SI (System International) unit.

In this juice label from Amsterdam you can see that energy (“energie”) is given in kilojoules and kilocalories.

To convert between calories and joules keep in mind:


1 calorie (cal) = 4.184 joules (J)

or

1 Calorie (kcal) = 1000 calories (cal) = 4184 joules (J) = 4.184 kilojoules (kJ)

or use this chart:

(you will need to **memorize** one of these conversion methods)



Specific Heat Capacity

Specific heat capacity (C) is defined as the amount of energy required to raise the temperature of one gram of a substance one degree Celsius. The unit for specific heat capacity is J/g°C. Specific heat capacity is sometimes referred to as just specific heat. Specific heat capacity values vary for different substances. It is a substance's specific heat that makes some substances feel cooler than others, even though both are at the same temperature. This phenomenon is due to the fact that some substances require little heat in order for their temperature to rise 1°C. Thus, substances with low specific heat values are said to be good conductors of heat. Most metals are considered to be good conductors because of their low C values. Substances with high C values are called insulators because they require large amounts of heat in order to warm up just a little. It is because of this, large bodies of water take a long time to warm in the summer and a long time to freeze in the winter. The chart below lists the specific heat capacities for some various substances.

Substance	Specific Heat Capacity (J/g°C)
Water (l)	4.184
Water (s)	2.09
Water (g)	2.01
Aluminum (s)	0.89
Iron (s)	0.45
Mercury (l)	0.14
Carbon (s)	0.71
Silver (s)	0.24
Gold (s)	0.13
Copper (s)	0.385

The specific heat capacity of a substance is needed when determining how much energy is absorbed or released when heat flows from one substance to another. The formula used is:

$$q = mC\Delta T$$

The Δ is the delta symbol and means "change in".

q – heat, measured in joules

m – the mass of a substance, measured in grams

C – the specific heat capacity of the substance that is being heated or cooled, measured in J/g°C

(-28°C) – the change in temperature (final temperature – initial temperature), measured in °C

Since you will either be given the initial and final temperatures or asked to find one of them, it may be better to express the formula as:

$$q = \text{mass} \times C \times (T_{\text{final}} - T_{\text{initial}})$$

If **q** is **positive**, it is an **endothermic change**, if **q** is **negative**, it is an **exothermic change**.

Example: How much heat energy is released to your body when a cup of hot tea containing 200.0 grams of water is cooled from 65°C to body temperature, 37°C?

$$q = \text{mass} \times C \times (T_{\text{final}} - T_{\text{initial}})$$

$$q = (200.0\text{g})(4.184 \text{ J/g } ^\circ\text{C})(37^\circ\text{C} - 65^\circ\text{C})$$

$$q = (200.0\text{g})(4.184 \text{ J/g } ^\circ\text{C})(-28^\circ\text{C})$$

$$q = -23000 \text{ J} = -23 \text{ kJ} = -5.6 \text{ kcal}$$

exothermic

Which C value of water should you use?

Water boils at 100°C and freezes at 0°C. If a temperature range is below 0°C, use the C value for solid water. If the temperature range is above 100°C use the C value for water vapor (g). If the temperature range is between 0°C and 100°C, use the C value for liquid water.

Phase Changes

Changes of state are physical changes where heat is absorbed or released. If a substance absorbs heat, the process is endothermic and it will change from a lower energy state to a higher energy state. If a substance releases heat, the process is exothermic and it will change from a higher energy state to a lower energy state. The lowest energy state is solid and the highest energy state is gas. The liquid state is intermediate between solid and gas.

As a substance melts, freezes, boils or condenses, heat is either absorbed or released. In molecular (covalent) compounds bonds between molecules are broken or formed; in ionic compounds bonds between ions are broken or formed. As the bonds between molecules, ions or atoms break, the particles move further and further apart; when bonds form particles move closer and closer together.

During physical changes of state, **there is no change in temperature**. Thus all energy being added is used to break bonds and any energy released is from the formation of bonds. For each change of state there is a term used to describe the process and an energy value associated with it.

Name	Change	Endo/Exo
Melting	Solid to Liquid	Endo
Freezing	Liquid to Solid	Exo
Sublimation	Solid to Gas	Endo
Deposition	Gas to Solid	Exo
Vaporization	Liquid to Gas	Endo
Condensation	Gas to Liquid	Exo

The chart below summarizes these processes:

Process	Name	Symbol	Endo/Exo	Bonds	Value for Water
Melting (s→l)	Heat of fusion	H_{fus}	Endo	Broken	335 J/g or 80. cal/g
Freezing (l→s)	Heat of solidification	H_{sol}	Exo	Formed	-335 J/g or -80. cal/g
Vaporization (l→g)	Heat of vaporization	H_{vap}	Endo	Broken	2259.0 J/g or 540. cal/g
Condensation (g→l)	Heat of condensation	H_{con}	Exo	Formed	-2259.0 J/g or -540. cal/g

*Each of the values listed above represent the amount of energy needed or released to (melt/freeze/vaporize/condense) **one gram** of water.

Because the temperature is constant, the formula: $q = mC\Delta T$ would not be accurate. Without a temperature change, $\Delta T = 0$ and our calculation of the heat absorbed or released would be zero. Since we know heat is being added or released a different formula must be used. At the melting/freezing point the formula to be used is: $q = H_x \times \text{mass (or moles)}$.

Example: Calculate the amount of energy needed to melt 5.00 grams of ice at 0°C .

$$q = H_{\text{fus}} \times \text{mass}$$

$$q = 335.0 \times 5.00$$

$$q = 1680 \text{ joules} \div 1000 = 1.68 \text{ kJ} \div 4.184 = 0.402 \text{ kcal}$$

The reverse process (freezing) is an exothermic process. Thus, instead of H_{fus} , H_{solid} (heat of solidification) is used. It has the same numeric value as a substance's H_{fus} but it is negative ($H_{\text{solid}}(\text{H}_2\text{O}) = -335.0 \text{ J/g}$). If the reverse process (freezing) were to occur above, then $q = -1680 \text{ J}$.

Example: Calculate the amount of energy needed to change 30.00 grams of water at 100°C to steam at 100°C .

$$q = H_{\text{vap}} \times \text{mass}$$

$$q = 2259.0 \times 30.00$$

$$q = 67770 \text{ joules} \div 1000 = 67.77 \text{ kJ} \div 4.184 = 16.20 \text{ kcal}$$

The reverse process (condensing) is an exothermic process. Thus, instead of H_{vap} , H_{cond} (heat of condensation) is used. It has the same numeric value as a substance's H_{vap} but it is negative ($H_{\text{cond}}(\text{H}_2\text{O}) = -2259.0 \text{ J/g}$). If the reverse process (condensing) were to occur above, then $q = -67770 \text{ J}$.

Note: The H_{fus} , H_{cond} , H_{vap} & H_{solid} values given above are in joules/gram. On occasion, however, you will be given values in joules/mole or kilojoules/mole. If this is the case, use the formulas: $q = \Delta H_{\text{fus}} \times \text{moles}$ or $q = \Delta H_{\text{vap}} \times \text{moles}$. **Be sure to convert mass → moles if necessary.** See the example below.

Example: Calculate the kilojoules of energy needed to melt 10.0 grams of NH_3 . $H_{\text{fus}} = 5.65 \text{ kJ/mole}$

First: Since the H_{fus} value is given in kJ/mole, I will use the formula $q = \Delta H_{\text{fus}} \times \text{moles}$.

Second: Convert mass to moles by dividing the given value by the gram formula mass of the substance.

$$10.0 \div 17.0 = 0.588 \text{ moles}$$

Third: Solve it.

$$q = \Delta H_{\text{fus}} \times \text{moles}$$

$$q = 5.65 \times 0.588$$

$$q = 3.32 \text{ kJ} \div 4.184 = 0.793 \text{ kcal}$$

Homework:

Part 1: Solve each specific heat capacity problem in kilojoules and Calories (kilocalories) or degrees Celsius.


Indicate if an endothermic or exothermic change takes place.

- If you fill a bathtub with 200. kg (convert kilograms to grams!) of water at 44°C , how much heat energy is lost as the water cools to a temperature of 21°C ?

2. On a cold winter day with a temperature of 4°C , you pick up a penny from the ground and put it in your pocket. If the penny has a mass of 1.85 grams, how much heat energy must be transferred to the coin to warm it to your body temperature, 37°C ?
3. Predict the final temperature of 2.50 kg (convert kilograms to grams) of water in a calorimeter if the water is at 25.0°C before 0.5 oz. of noodles, which contain 54 Calories (convert to joules), are burned?
4. How much heat is lost by 4.0 L (convert L to mL & convert mL to grams; density of water 1.0 g/mL) of water that is cooled from 87°C to 21°C ?
5. If 980 kJ of energy are added to 6.2 kg of water at 18°C , what is the final temperature of the water?
6. How much heat energy is needed to raise the temperature of a 425.0 gram aluminum baking sheet from room temperature, 25°C , to a baking temperature of $200.^{\circ}\text{C}$?
7. Calculate the amount of energy needed to raise the temperature of 100. grams of gold from $30.^{\circ}\text{C}$ to $40.^{\circ}\text{C}$.
8. Calculate the amount of energy released as 225 grams of mercury cools from $100.^{\circ}\text{C}$ to 27°C .
9. Which of the following would be the best choice as an insulator, gold, iron, aluminum or water? Explain
10. Which of the following is the best conductor of heat, carbon, water, copper, or mercury? Explain.
11. If a q value is positive, is a reaction endothermic or exothermic?

Part 2: Use the diagram to the bottom right to identify the process occurring.

1. _____
2. _____
3. _____
4. _____
5. _____
6. _____



Part 3: Solve each phase change problems in kilojoules and Calories (kilocalories). Indicate if an endothermic or exothermic change takes place.

1. Calculate the amount of energy required to change 15.0 grams of ice at 0°C to 15.0 grams of water at 0°C . $H_{\text{fus}} = 80.0$ cal/gram

2. If the heat of fusion of water is 340 J/g, calculate the amount of heat energy required to change 40.0 grams of ice at 0°C to 40.0 grams of water at 0°C .

3. Calculate the amount of energy released as 100.0 grams of steam are condensed into liquid water at 100°C .

4. Calculate the amount of energy released as 35.0 grams of water at 0°C changes to 35.0 grams of ice at 0°C .

5. Calculate the amount of heat energy released as 220.0 grams of water at 0°C is changed to 220.0 grams of ice at 0°C .

6. Calculate the amount of energy absorbed as 400.0 grams of water is vaporized at 100°C .

7. If the heat of fusion of FeO is 32.2 kJ/mol, calculate the amount of heat energy required to melt 17.7 grams of FeO.

8. Calculate the kilojoules of energy needed to vaporize 70.0 grams of NH_3 . $H_{\text{vap}} = 23.4 \text{ kJ/mol}$
9. Calculate the amount of energy released as 55.0 grams of Br_2 is changed from liquid to solid. $H_{\text{fus}} = 10.72 \text{ kJ/mol}$
10. The energy required to melt a solid into a liquid is called —
11. The energy released when freezing a liquid to a solid is called —
12. The energy released when condensing a gas to a liquid is called —
13. The energy required to vaporize a liquid into a gas is called —
14. Heat of condensation is an (endothermic / exothermic) process. Circle one.
15. Heat of fusion is an (endothermic / exothermic) process. Circle one.
16. Heat of vaporization is an (endothermic / exothermic) process. Circle one.
17. Heat of solidification is an (endothermic / exothermic) process. Circle one.
18. Heat of fusion is used at the: (melting point / boiling point / freezing point / condensing point) Circle one.
19. Heat of vaporization is used at the: (melting point / boiling point / freezing point / condensing point) Circle one.
20. Heat of solidification is used at the: (melting point / boiling point / freezing point / condensing point) Circle one.
21. Heat of condensation is used at the: (melting point / boiling point / freezing point / condensing point) Circle one.

Part 4: Multiple Choice Problems - Choose the best answer for each of the following.

1. How much heat is required to raise the temperature of 100. g of Fe_2O_3 from 5.0°C to 25.0°C ? (Specific heat Fe_2O_3 , $0.634 \text{ J}\cdot\text{g}^{-1}\cdot^\circ\text{C}^{-1}$)
- (A) 1.58 kJ (B) 1.27 kJ (C) 0.845 kJ (D) 0.0634 kJ (E) 1.902 kJ

2. The specific heats of several metals are given in the table. If the same number of Joules were applied to the same mass of each metal, which metal would show the greatest temperature change?

Substance	Specific Heat, $\text{J}\cdot\text{g}^{-1}\cdot^\circ\text{C}^{-1}$
Al	0.900
Au	0.129
Cu	0.385
Hg	0.139

- (A) Al (B) Au (C) Cu (D) Hg (E) all would be equal