


Chapter 4 Collected AP Exam Free Response Questions 1980 - 2009

1981 - #3a


A 1.2156-gram sample of a mixture of CaCO_3 and Na_2SO_4 was analyzed by dissolving the sample and completely precipitating the Ca^{2+} as CaC_2O_4 . The CaC_2O_4 was dissolved in sulfuric acid and the resulting $\text{H}_2\text{C}_2\text{O}_4$ was titrated with a standard KMnO_4 solution.

(a) Write the balanced equation for the titration reaction, shown balanced below.



Indicate which substance is the oxidizing agent and which substance is the reducing agent.

1983 - #8a & b



(a) Write the correctly balanced half-reaction and net ionic equation for the skeletal equation shown above.

(b) Identify the oxidizing agent and the reducing agent in this reaction.

1987 - #7b & d

In 1884 the Swedish chemist Svante Arrhenius proposed that salts dissociate into two or more separate, independent, ionic fragments when they dissolve in water.

(b) Give one piece of experimental evidence that the particles formed when a salt dissolves in water are charged.

(d) Explain why hydrogen chloride, HCl , dissociates when it dissolves in water, but not when it dissolves in benzene.

1998 - #1a

Solve the following problem related to the solubility equilibria of some metal hydroxides in aqueous solution.


(a) The solubility of $\text{Cu}(\text{OH})_2$ is 1.72×10^{-6} gram per 100. milliliters of solution at 25°C .

(i) Write the balanced chemical equation for the dissociation of $\text{Cu}(\text{OH})_2(\text{s})$ in aqueous solution.

(ii) Calculate the solubility (in moles per liter) of $\text{Cu}(\text{OH})_2$ at 25°C .

2000 - #3ci

A 0.345 g sample of anhydrous BeC_2O_4 , which contains an inert impurity, was dissolved in sufficient water to produce 100. mL of solution. A 20.0 mL portion of the solution was titrated with $\text{KMnO}_4(\text{aq})$. The balanced equation for the reaction that occurred is as follows.



(i) Identify the reducing agent in the titration reaction.

2001 - #1a

Answer the following questions relating to the solubility of the chlorides of silver and lead.

(a) At 10°C , 8.9×10^{-3} g of $\text{AgCl}(\text{s})$ will dissolve in 100. mL of water.

(i) Write the equation for the dissociation of $\text{AgCl}(\text{s})$ in water.

(ii) Calculate the solubility, in mol L^{-1} , of $\text{AgCl}(\text{s})$ in water at 10°C .

2001 - #5c & e

Solution 1



0.10 M
 $\text{Pb}(\text{NO}_3)_2$

Solution 2



0.10 M
 NaCl

Solution 3



0.10 M
 KMnO_4

Solution 4



0.10 M
 $\text{C}_2\text{H}_5\text{OH}$

Solution 5




0.10 M
 $\text{KC}_2\text{H}_3\text{O}_2$

Answer the questions below that relate to the five aqueous solutions at 25°C shown above.

- (c) Identify a pair of the solutions that would produce a precipitate when mixed together. Write the formula of the precipitate.
- (e) Which solution would be the least effective conductor of electricity? Explain.

2002 - #2a

Answer parts (a) through (e) below, which relate to reactions involving silver ion, Ag^+ . The reaction between silver ion and solid zinc is represented by the following equation.



- (a) A 1.50 g sample of Zn is combined with 250. mL of 0.110 M AgNO_3 at 25°C.
- (i) Identify the limiting reactant. Show calculations to support your answer.
- (ii) On the basis of the limiting reactant that you identified in part (i), determine the value of $[\text{Zn}^{2+}]$ after the reaction is complete. Assume that volume change is negligible.


2002B - #3a

Nitrogen monoxide, $\text{NO}(g)$, and carbon monoxide, $\text{CO}(g)$, are air pollutants generated by automobiles. It has been proposed that under suitable conditions these two gases could react to form $\text{N}_2(g)$ and $\text{CO}_2(g)$, which are components of unpolluted air.

- (a) Write a balanced equation for the reaction described above. Indicate whether the carbon in CO is oxidized or whether it is reduced in the reaction. Justify your answer.

2003B - #2b


In a reaction vessel, 0.600 mol of $\text{Ba}(\text{NO}_3)_2(s)$ and 0.300 mol of $\text{H}_3\text{PO}_4(aq)$ are combined with deionized water to a final volume of 2.00 L. The reaction represented below occurs.



- (i) Calculate the mass of $\text{Ba}_3(\text{PO}_4)_2(s)$ formed.
- (iii) What is the concentration, in mol L^{-1} , of the nitrate ion, $\text{NO}_3^-(aq)$, after the reaction reaches completion?

2003B - #5a

Oxalic acid, $\text{H}_2\text{C}_2\text{O}_4$, is a primary standard used to determine the concentration of potassium permanganate, KMnO_4 , in solution. The equation for the reaction is as follows.



A student dissolves a sample of oxalic acid in a flask with 30 mL of water and 2.00 mL of 3.00 M H_2SO_4 . The KMnO_4 solution of unknown concentration is in a 25.0 mL buret. In the titration, the KMnO_4 solution is added to the solution containing oxalic acid.

- (a) What chemical species is being oxidized in the reaction?

2004 - #5

Each flask contains one of the following solutions:

$\text{Pb}(\text{NO}_3)_2$
 NaCl
 K_2CO_3



Each flask contains one of the following solutions:

AgNO_3
 BaCl_2




- (a) When the student combined a sample of solution Q with a sample of solution X, a precipitate formed. A precipitate also formed when samples of solutions Q and Y were combined.
- (i) Identify solution Q.
- (ii) Write the chemical formulas for each of the two precipitates.
- (b) When solution Q is mixed with solution R, a precipitate forms. However, no precipitate forms when solution Q is mixed with solution S.
- (i) Identify solution R and solution S.

- (ii) Write the chemical formula of the precipitate that forms when solution *Q* is mixed with solution *R*.
- (c) The identity of solution *X* and solution *Y* are to be determined using only the following solutions: 1.0 *M* Pb(NO₃)₂, 1.0 *M* NaCl, and 1.0 *M* K₂CO₃.
- (i) Describe a procedure to identify solution *X* and solution *Y*.
- (ii) Describe the observations that would allow you to distinguish between solution *X* and solution *Y*.
- (iii) Explain how the observations would enable you to distinguish between solution *X* and solution *Y*.

2006 - #1a

Answer the following questions that relate to solubility of salts of lead and barium.


A saturated solution is prepared by adding excess PbI₂(*s*) to distilled water to form 1.0 L of solution at 25°C. The concentration of Pb²⁺(*aq*) in the saturated solution is found to be 1.3 × 10⁻³ *M*. The chemical equation for the dissolution of PbI₂(*s*) in water is shown below.



- (ii) Calculate the molar concentration of I⁻(*aq*) in the solution.

2007 - #1c & d


HF(*aq*) reacts with NaOH(*aq*) according to the reaction represented below.



A volume of 15 mL of 0.40 *M* NaOH(*aq*) is added to 25 mL of 0.40 *M* HF(*aq*) solution. Assume that volumes are additive.

- (c) Calculate the number of moles of HF(*aq*) remaining in the solution.
- (d) Calculate the molar concentration of F⁻(*aq*) in the solution.

2007 - #5a & b




The mass percent of iron in a soluble iron(II) compound is measured using a titration based on the balanced equation above.


- (a) What is the oxidation number of manganese in the permanganate ion, MnO₄⁻(*aq*)?
- (b) Identify the reducing agent in the reaction represented above.

2008B - #3

A 0.150 g sample of solid lead(II) nitrate is added to 125 mL of 0.100 *M* sodium iodide solution. Assume no change in volume of the solution. The chemical reaction that takes place is represented by the following equation.




- (a) List an appropriate observation that provides evidence of a chemical reaction between the two compounds.
- (b) Calculate the number of moles of each reactant.
- (c) Identify the limiting reactant. Show calculations to support your identification.
- (d) Calculate the molar concentration of NO₃⁻(*aq*) in the mixture after the reaction is complete.
- (e) Circle the diagram below that best represents the results after the mixture reacts as completely as possible. Explain the reasoning used in making your choice.



2010B - #3

A sample of ore containing the mineral tellurite, TeO_2 , was dissolved in acid. The resulting solution was then reacted with a solution of $\text{K}_2\text{Cr}_2\text{O}_7$ to form telluric acid, H_2TeO_4 . The unbalanced chemical equation for the reaction is given below.



- Identify the molecule or ion that is being oxidized in the reaction.
- Give the oxidation number of Cr in the $\text{Cr}_2\text{O}_7^{2-}(aq)$ ion.
- Balance the chemical equation given above by writing the correct lowest whole-number coefficients on the dotted lines.