HW 11_4: Due 2/16/17 Write the letter of the correct answer on the line in front of the question.

1	What mass of KBr (molar mass 119 g mol ⁻¹) is required to make 250. mL of a 0.400 M KBr								
solution?									
(A) 0.595 g	(B) 1.19 g	(C) 2.50 g	(D) 11.9 g	(E) 47.6 g					
2	A sample of a solution	of RbCl (molar m	ass 121 o m	ol ⁻¹) contains 11.0 percent RbCl by mass	s				
From the following information, what is needed to determine the molarity of RbCl in the solution?									
I Mass of the sample									
	II. Volume of the sample								
	III. Temperature of the sa	mple							
(A) I only (B) II only (C) I and II only									
(D) II and III only (E) I, II, and III									
3.	Which of the following	ng aqueous solutio	ns has the h	ighest boiling point at 1.0 atm?					
(A) $0.20 M \text{ CaCl}_2$ (B) $0.25 M \text{ Na}_2\text{SO}_4$ (C) $0.30 M \text{ NaCl}$									
()	(D) $0.30 M \text{ KBr}$ (E) $0.40 M \text{ C}_6 \text{H}_{12} \text{O}_6$								
4 Molarity units are most appropriate in calculating which of the following?									
(A) freezing poi	nt depression	(C) boiling point elevation							
() - 81	(D) surface tensi	on	(1	E) osmotic pressure					
5.	The weight of H_2SO_4 (1)	molecular weight (98.1) in 500	0 milliliters of a 6.00-molar solution is					
(A) 3.10 grams	(B) 12.0 grams (C) 29.4	4 grams (D) 294	grams (I	E) 300. grams					
6.	Which of the following	solutions has the lo	owest boilin	g point?					
(A) $0.20 m C_6 H_1$	$\overline{O}_{12}O_{6}$, glucose (B) 0.20	$m \mathrm{NH}_4 \mathrm{Br}$	((C) 0.20 m ZnSO ₄					
	(D) $0.20 \ m \ \text{KMnO}_4$	(E) 0.20	$m \operatorname{MgCl}_2$						
7.	If the temperature of ar	n aqueous solution	of NaCl is	increased from 20 °C to 90 °C, which of					
the following sta	atements is true?	1							
(A) The density	of the solution remains unc	hanged.	(B) The m	olarity of the solution remains unchanged	d.				
(C) The molality of the solution remains unchanged. (D) The mole fraction of solute decreases.									
	(E) The	e mole fraction of s	solute increa	ases.					
8.	I. freezing point	depression	II. osmotic	pressure III. vapor pressure					
Mole fractions a	re typically used to calculat	te which properties	s for solutio	ns containing nonvolatile solutes?					
(A) I only	(B) II only (C) III o	only (D) I an	d II only	(E) II and III only					
9.	Fish kills are often obs	served in lakes and	l ponds in th	he summer but rarely in the winter. A					
contributing fact	tor is the use of oxygen by c	lecaying algae. A	nother facto	or is:					
(A) the higher so	olubility of toxic metals in t	he summer							
(B) the decreased solubility of oxygen at higher temperature									
(C) the high temperature itself kills the fish									
(D) the toxicity of decaying algae									
(E) soluble nutri	ents are generally less solul	ble at higher tempe	eratures						
10	Ethyl alcohol, C ₂ H ₅ C	H, and water beco	ome noticeal	bly warmer when mixed. This is due to:					
(A) the decrease in volume when they are mixed									
(B) smaller attractive forces in the mixture than in the pure liquids									
(C) the hydrogen bonding of the two liquids									
(D) the change in vapor pressure observed (F) stronger attractive forces in the mixture than in the pure liquids									
(E) subliger attra	active forces in the mixture	man in the pute fi	quius						

11	The molality of the glucose in a 1.0-molar glucose solution can be obtained by using which of							
the following?								
(A) Volume of the	ne solution	(B) Temperatu	(B) Temperature of the solution (C) Solubility of glucose in water					
	(D) Degree of dissociation of glucose		(E) Density of the solution					
12 What is the mole fraction of ethanol, C_2H_5OH , in an aqueous solution in which the ethanol concentration is 5.02 molal?								
(A) 0.0046	(B) 0.076	(C) 0.083	(D) 0.20	(E) 0.72				
13 If equal numbers of moles of each of the following are dissolved in 1 kg of distilled water, the one with the lowest boiling point will be:(A) NaF(B) AlCl ₃ (C) Mg(C ₂ H ₃ O ₂) ₂ (D) CH ₃ CH ₂ COOH(E) C ₆ H ₆								
 14. A solution of hydrochloric acid has a density of 1.15 grams per mL and is 30.% by weight HCl. (a) What is the molarity of this solution of HCl? (b) What volume of this solution should be taken in order to prepare 5.0 liters of 0.20-molar hydrochloric acid by dilution with water? (a) In order to obtain a pracise concentration, the 0.20 molar hydrochloric acid is standardized against pure HgO 								
(molecular weight = 216.59) by titrating the OH ² produced according to the following quantitative reaction:								

$$\operatorname{HgO}(s) + 4I^{-} + \operatorname{H_2O} \rightarrow \operatorname{HgI_4^{2-}} + 2O$$

In a typical experiment, 0.7147 gram of HgO required 31.67 milliliters of the hydrochloric acid solution for titration. Based on these data, what is the molarity of the HCl solution expressed to four significant figures?